



TITLE:

Analytic smoothing effect for nonlinear
Schrodinger equations with quadratic
interaction (Harmonic Analysis and
Nonlinear Partial Differential Equations)

AUTHOR(S):

HOSHINO, Gaku; OZAWA, Tohru

CITATION:

HOSHINO, Gaku ...[et al]. Analytic smoothing effect for nonlinear Schrodinger equations with quadratic interaction (Harmonic Analysis and Nonlinear Partial Differential Equations). 数理解析研究所講究録別冊 2014, B49: 1-12: KJ00009529619.

ISSUE DATE:

2014-06

URL:

<http://hdl.handle.net/2433/226239>

RIGHT:

c 2014 by the Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

Analytic smoothing effect for nonlinear Schrödinger equations with quadratic interaction

By

Gaku HOSHINO* and Tohru OZAWA**

Abstract

We prove the global existence of analytic solutions to a system of nonlinear Schrödinger equations with quadratic interaction in space dimensions $n \geq 3$, under the mass resonance condition $M = 2m$ for sufficiently small Cauchy data with exponential decay. For $n \geq 4$ the smallness assumption on the data is imposed in terms of the critical Sobolev space $\dot{H}^{n/2-2}$.

§ 1. Introduction

We consider the Cauchy problem for a system of nonlinear Schrödinger equations of the form

$$(1.1) \quad \begin{cases} i\partial_t u + \frac{1}{2m}\Delta u = \lambda v\bar{u}, \\ i\partial_t v + \frac{1}{2M}\Delta v = \mu u^2 \end{cases}$$

in space dimension $n \geq 3$, under the mass resonance condition $M = 2m$, where u, v are complex valued functions of $(t, x) \in \mathbb{R} \times \mathbb{R}^n$, Δ is the Laplacian in \mathbb{R}^n , and $\lambda, \mu \in \mathbb{C}$.

The Cauchy problem for (1.1) has been studied in [11, 12, 13, 16, 20]. Particularly, Lagrangian formalism was described in [20], the existence of ground states was proved in [16], and analytic solutions were studied in [20] in terms of the generator of Galilei transforms. In this paper, we give another framework of analytic smoothing effect for (1.1).

Received August 26, 2013. Revised December 4, 2013.

2010 Mathematics Subject Classification(s): 35Q55

Key Words: Nonlinear Schrödinger equations, Analyticity

Supported by Grants for Excellent Graduate Schools, MEXT, Japan.

*Department of Applied Physics, Waseda University, Tokyo 169-8555, Japan.

e-mail: gaku-hoshino@ruri.waseda.jp

**Department of Applied Physics, Waseda University, Tokyo 169-8555, Japan.

e-mail: txozawa@waseda.jp

To state our main result precisely, we introduce the following notation. For $1 \leq p \leq \infty$, $L^p = L^p(\mathbb{R}^n)$ is the usual Lebesgue space. For $s \in \mathbb{R}$, $H_p^s = (1 - \Delta)^{-s/2} L^p$ and $\dot{H}_p^s = (-\Delta)^{-s/2} L^p$ are the Sobolev space and the homogeneous Sobolev space, respectively. $\dot{L}_s^2 = |x|^{-s} L^2$ is the weighted Lebesgue space. For $1 \leq p, q \leq \infty$, $\dot{B}_{p,q}^s = \dot{B}_{p,q}^s(\mathbb{R}^n)$ is the homogeneous Besov space and $L^{p,q} = L^{p,q}(\mathbb{R})$ is the Lorentz space with second exponent q . We put $\dot{B}_{p,2}^s = \dot{B}_p^s$. For $t \in \mathbb{R}$, $U_m(t) = e^{i\frac{t}{2m}\Delta}$ is the free propagator with mass m and $J_m(t) = x + i\frac{t}{m}\nabla = U_m(t)xU_m(-t)$ is the generator of Galilei transforms. For any $t \neq 0$, $J_m(t)$ is represented as $J_m(t) = M_m(t)i\frac{t}{m}\nabla M_m(-t)$, where $M_m(t) = e^{i\frac{m}{2t}|x|^2}$. \mathcal{F} denotes the Fourier transform defined by $(\mathcal{F}f)(\xi) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-i\xi \cdot x} f(x) dx$. For $k > 0$, $\delta \in \mathbb{R}^n$ and $t \in \mathbb{R}$ we define the following operator [9, 15, 21, 27]:

$$P_{k\delta,m}(t) = U_m(t)e^{k\delta \cdot x}U_m(-t),$$

where \cdot is the usual scalar product in \mathbb{R}^n . For any $t \neq 0$, $P_{k\delta,m}(t)$ is represented as

$$P_{k\delta,m}(t) = M_m(t)\mathcal{F}^{-1}e^{-t\frac{k\delta}{m} \cdot \xi}\mathcal{F}M_m(-t).$$

Let \mathcal{B} be a suitable Banach space consisting of functions on space-time $\mathbb{R} \times \mathbb{R}^n$. Let D be an open subset of \mathbb{R}^n . We introduce the following function spaces:

$$\begin{aligned} G_k^D(x; \dot{H}_2^s) &= \left\{ \phi \in \dot{H}_2^s ; \|\phi; G_k^D(x; \dot{H}_2^s)\| \equiv \sup_{\delta \in D} \|e^{k\delta \cdot x} \phi; \dot{H}_2^s\| < \infty \right\}, \\ G_k^D(x; L^2 \cap \dot{L}_s^2) &= \left\{ \phi \in L^2 \cap \dot{L}_s^2 ; \|\phi; G_k^D(x; L^2 \cap \dot{L}_s^2)\| \equiv \sup_{\delta \in D} \|e^{k\delta \cdot x} \phi; L^2 \cap \dot{L}_s^2\| < \infty \right\}, \\ G_k^D(J_m; \mathcal{B}) &= \left\{ u \in \mathcal{B} ; \|u; G_k^D(J_m; \mathcal{B})\| \equiv \sup_{\delta \in D} \|P_{k\delta,m}u; \mathcal{B}\| < \infty \right\}. \end{aligned}$$

Our main purpose in this paper is twofold. One is to study analytic smoothing effect for nonlinear Schrödinger equations in terms of Hardy spaces of analytic functions on tube domains, which provides a wider framework than [21]. The other is to apply this framework to (1.1) and to improve the result in [20]. The space $G_k^D(J_m; \mathcal{B})$ is characterized by convergence of power series which respect to $k\delta \cdot J_m$ in \mathcal{B} ([21]). The Cauchy problem for nonlinear Schrödinger equations in the function space in terms of absolutely convergence of power series with respect to $k\delta \cdot J_m$ is studied by ([19, 20, 33, 34]).

For $n \geq 4$ we introduce the following function spaces:

$$\begin{aligned} \mathcal{X}_0 &= L^\infty(\mathbb{R}; L^2) \cap L^2(\mathbb{R}; L^{2*}), \\ \dot{\mathcal{X}} &= L^\infty(\mathbb{R}; \dot{H}_2^{n/2-2}) \cap L^2(\mathbb{R}; \dot{B}_{2*}^{n/2-2}) \end{aligned}$$

with associated norm

$$\begin{aligned}\|u; \mathcal{X}_0\| &= \|u; L^\infty(\mathbb{R}; L^2)\| + \|u; L^2(\mathbb{R}; L^{2^*})\|, \\ \|u; \dot{\mathcal{X}}\| &= \|u; L^\infty(\mathbb{R}; \dot{H}_2^{n/2-2})\| + \|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\|,\end{aligned}$$

where $2^* = 2n/(n-2)$ is the Sobolev critical exponent. For $n = 3$ we introduce the following function space:

$$\mathcal{Y}_0 = L^\infty(\mathbb{R}; L^2) \cap L^{4,2}(\mathbb{R}; L^3)$$

with associated norm

$$\|u; \mathcal{Y}_0\| = \|u; L^\infty(\mathbb{R}; L^2)\| + \|u; L^{4,2}(\mathbb{R}; L^3)\|.$$

We define $|J_m|^s$ by

$$|J_m|^s(t) = U_m(t)|x|^s U_m(-t)$$

for $t \in \mathbb{R}$. For any $t \neq 0$, $|J_m|^s(t)$ is represented as [14]

$$|J_m|^s(t) = M_m(t) \left(-\frac{t^2}{m^2} \Delta \right)^{s/2} M_m(-t).$$

We put

$$\dot{\mathcal{Y}}_m = |J_m|^{-1/2} \mathcal{Y}_0.$$

We treat (1.1) in the following function spaces:

$$\begin{aligned}G_k^D(J_m) &= \mathcal{X}_0 \cap \dot{G}_k^D(J_m) && \text{for } n \geq 4, \\ G_k^D(J_m) &= G_{k,0}^D(J_m) \cap \dot{G}_k^D(J_m) && \text{for } n = 3,\end{aligned}$$

where

$$\begin{aligned}\dot{G}_k^D(J_m) &= G_k^D(J_m; \dot{H}_2^{n/2-2}) && \text{for } n \geq 4, \\ G_{k,0}^D(J_m) &= G_k^D(J_m; \mathcal{Y}_0), \quad \dot{G}_k^D(J_m) = G_k^D(J_m; \dot{\mathcal{Y}}_m) && \text{for } n = 3.\end{aligned}$$

We define $B_k^D(\rho)$ as follows

$$\begin{aligned}B_k^D(\rho) &= \left\{ \phi \in G_k^D(x; \dot{H}_2^{n/2-2}) ; \sup_{\delta \in D} \|e^{\delta \cdot x} \phi; \dot{H}_2^{n/2-2}\| \leq \rho \right\} && \text{for } n \geq 4, \\ B_k^D(\rho) &= \left\{ \phi \in G_k^D(x; L^2 \cap \dot{L}_{1/2}^2) ; \sup_{\delta \in D} \|e^{\delta \cdot x} \phi; \dot{L}_{1/2}^2\| \leq \rho \right\} && \text{for } n = 3.\end{aligned}$$

We consider the following integral equations of (1.1) with the Cauchy data (ϕ, ψ) at $t = 0$:

$$\begin{cases} u(t) = U_m(t)\phi - i \int_0^t U_m(t-s)\lambda(v\bar{u})(s)ds, \\ v(t) = U_{2m}(t)\psi - i \int_0^t U_{2m}(t-s)\mu u^2(s)ds. \end{cases}$$

We state our main result.

Theorem 1.1. *Let $n \geq 3$. There exists $\rho > 0$ such that for any domain $D \subset \mathbb{R}^n$ satisfying $0 \in D$, $-D = D$ and any $(\phi, \psi) \in B_1^D(\rho) \times B_2^D(\rho)$ (1.1) has a unique pair of solutions $(u, v) \in G_1^D(J_m) \times G_2^D(J_{2m})$.*

Remark 1. Under the assumptions of Theorem 1.1, if $\phi \in \dot{H}_2^{n/2-2}$ satisfies $\sup_{\delta \in D} \|e^{k\delta \cdot x} \phi; \dot{H}_2^{n/2-2}\| < \infty$, then $\phi \in L^1 \cap L^{n/2}$ ([21, 40]).

Remark 2. Under the assumptions of Theorem 1.1, let $u \in G_k^D(J_{lm})$, $l > 0$. Then for any $t \neq 0$, $u(t)$ is real analytic and has an analytic continuation to $\mathbb{R}^n + i\frac{t}{lm}kD$ ([17, 18, 21, 27, 35, 40]).

§ 2. Preliminaries

In this section we collect some basic lemmas. We define the following integral operator

$$(\Xi_m f)(t) = \int_0^t U_m(t-s)f(s)ds.$$

For $n \geq 3$, we say that (q, r) is an admissible pair if

$$\frac{2}{q} = n\left(\frac{1}{2} - \frac{1}{r}\right),$$

where $r \in [2, \frac{2n}{n-2}]$.

Lemma 2.1. ([3, 22, 38, 42, 44]) *Let $n \geq 1$, $s \in \mathbb{R}$ and let (q, r) and (q_j, r_j) be admissible pairs, $j = 1, 2$. Then the following estimates hold:*

$$\begin{aligned} \|U_m(\cdot)\phi; L^q(\mathbb{R}; L^r)\| &\leq C\|\phi; L^2\|, \\ \|\Xi_m f; L^{q_2}(\mathbb{R}; L^{r_2})\| &\leq C\|f; L^{q'_1}(\mathbb{R}; L^{r'_1})\|, \\ \|U_m(\cdot)\phi; L^q(\mathbb{R}; \dot{B}_r^s)\| &\leq C\|\phi; \dot{H}_2^s\|, \\ \|\Xi_m f; L^{q_2}(\mathbb{R}; \dot{B}_{r_2}^s)\| &\leq C\|f; L^{q'_1}(\mathbb{R}; \dot{B}_{r'_1}^s)\|, \end{aligned}$$

where p' is the dual exponent to p defined by $1/p + 1/p' = 1$.

Lemma 2.2. ([30, 31]) *Let $n \geq 1$ and let (q, r) and (q_j, r_j) be admissible pairs, $j = 1, 2$. Then the following estimates hold:*

$$\begin{aligned} \|U_m(\cdot)\phi; L^{q,2}(\mathbb{R}; L^r)\| &\leq C\|\phi; L^2\|, \\ \|\Xi_m f; L^\infty(\mathbb{R}; L^2)\| + \|\Xi_m f; L^{q_2,2}(\mathbb{R}; L^{r_2})\| &\leq C\|f; L^{q'_1,2}(\mathbb{R}; L^{r'_1})\|. \end{aligned}$$

Lemma 2.3. ([2, 26]) *Let $n \geq 1$. Let $1 \leq p, q \leq \infty, 1/p_1 + 1/p_2 = 1/p$ and $s > 0$. Then there exists $C > 0$ such that the following estimate*

$$\|uv; \dot{B}_{p,q}^s\| \leq C\left(\|u; \dot{B}_{p_1,q}^s\|\|v; L^{p_2}\| + \|u; L^{p_2}\|\|v; \dot{B}_{p_1,q}^s\|\right)$$

holds.

Lemma 2.4. ([13]) *Let $n = 3$. Then there exists $C > 0$ such that the following estimates hold:*

$$\begin{aligned} \|uv; L^{3/2}\| &\leq C|t|^{-1/2}\left\|\left|J_m\right|^{1/2}(t)u; L^2\right\|\|v; L^3\|, \\ \left\|\left|J_m\right|^{1/2}(t)(v\bar{u}); L^{3/2}\right\| &\leq C|t|^{-1/2}\left(\left\|\left|J_m\right|^{1/2}(t)u; L^2\right\|\left\|\left|J_{2m}\right|^{1/2}(t)v; L^3\right\| \right. \\ &\quad \left. + \left\|\left|J_m\right|^{1/2}(t)u; L^3\right\|\left\|\left|J_{2m}\right|^{1/2}(t)v; L^2\right\|\right), \\ \left\|\left|J_{2m}\right|^{1/2}(t)u^2; L^{3/2}\right\| &\leq C|t|^{-1/2}\left\|\left|J_m\right|^{1/2}(t)u; L^2\right\|\left\|\left|J_m\right|^{1/2}(t)u; L^3\right\| \end{aligned}$$

for all $t \neq 0$ and $m > 0$.

§ 3. Proof of Theorem 1.1.

Let $(\phi, \psi) \in B_1^D(\rho) \times B_2^D(\rho)$. We consider the mapping $(\Phi, \Psi) : (u, v) \mapsto (\Phi(u, v), \Psi(u, v))$ defined by

$$\begin{cases} \Phi(u, v)(t) = U_m(t)\phi - i\lambda\Xi_m(v\bar{u})(t), \\ \Psi(u, v)(t) = U_{2m}(t)\psi - i\mu\Xi_{2m}(u^2)(t) \end{cases}$$

on $G_1^D(J_m) \times G_2^D(J_{2m})$. For $R, \varepsilon > 0$ we define the metric spaces:

$$\begin{aligned} X(R, \varepsilon) &= \left\{ (u, v) \in G_1^D(J_m) \times G_2^D(J_{2m}) ; \right. \\ &\quad \left. \|(u, v); \mathcal{X}_0 \times \mathcal{X}_0\| \leq R, \|(u, v); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m})\| \leq \varepsilon \right\} \quad \text{for } n \geq 4, \end{aligned}$$

$$\begin{aligned} X(R, \varepsilon) &= \left\{ (u, v) \in G_1^D(J_m) \times G_2^D(J_{2m}) ; \right. \\ &\quad \left. \|(u, v); G_{1,0}^D(J_m) \times G_{2,0}^D(J_{2m})\| \leq R, \|(u, v); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m})\| \leq \varepsilon \right\} \quad \text{for } n = 3 \end{aligned}$$

with metric

$$d((u, v), (u', v')) = \|(u - u', v - v'); G_1^D(J_m) \times G_2^D(J_{2m})\|.$$

We see that $(X(R, \varepsilon), d)$ is a complete metric space.

Let $(u, v) \in G_1^D(J_m) \times G_2^D(J_{2m})$. We have

$$\begin{aligned} P_{\delta, m}\Phi(u, v) &= U_m e^{\delta \cdot x} U_m^{-1} \Phi(u, v) \\ &= U_m e^{\delta \cdot x} \phi - i\lambda \Xi_m \left(U_m e^{\delta \cdot x} U_m^{-1} (v\bar{u}) \right) \\ &= U_m e^{\delta \cdot x} \phi - i\lambda \Xi_m \left((U_{2m} e^{2\delta \cdot x} U_{2m}^{-1} v) \overline{(U_m e^{-\delta \cdot x} U_m^{-1} u)} \right) \\ &= U_m e^{\delta \cdot x} \phi - i\lambda \Xi_m \left((P_{2\delta, 2m} v) (\overline{P_{-\delta, m} u}) \right), \\ P_{2\delta, 2m}\Psi(u, v) &= U_{2m} e^{2\delta \cdot x} U_{2m}^{-1} \Psi(u, v) \\ &= U_{2m} e^{2\delta \cdot x} \psi - i\mu \Xi_{2m} \left(U_{2m} e^{2\delta \cdot x} U_{2m}^{-1} u^2 \right) \\ &= U_{2m} e^{2\delta \cdot x} \psi - i\mu \Xi_{2m} \left((U_m e^{\delta \cdot x} U_m u)^2 \right) \\ &= U_{2m} e^{2\delta \cdot x} \psi - i\mu \Xi_{2m} \left((P_{\delta, m} u)^2 \right). \end{aligned}$$

For $n \geq 4$, by Lemmas 2.1 and 2.4, the Hölder's inequality with $1/2 = 1/2^* + 1/n$, and the Sobolev embeddings $\dot{B}_{2^*}^{n/2-2} \subset \dot{H}_{2^*}^{n/2-2} \subset L^n$, we estimate

$$\begin{aligned} \|\Phi(u, v); \mathcal{X}_0\| &\leq C\|\phi; L^2\| + C\|v\bar{u}; L^1(\mathbb{R}; L^2)\| \\ &\leq C\|\phi; L^2\| + C\|v; L^2(\mathbb{R}; L^n)\| \|u; L^2(\mathbb{R}; L^{2^*})\| \\ &\leq C\|\phi; L^2\| + C\|v; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \|u; L^2(\mathbb{R}; L^{2^*})\|, \\ \|\Phi(u, v); \dot{\mathcal{X}}\| &\leq C\|\phi; \dot{H}_2^{n/2-2}\| + C\|v\bar{u}; L^1(\mathbb{R}; \dot{B}_2^{n/2-2})\| \\ &\leq C\|\phi; \dot{H}_2^{n/2-2}\| + C\left(\|v; L^2(\mathbb{R}; L^n)\| \|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \right. \\ &\quad \left. + \|u; L^2(\mathbb{R}; L^n)\| \|v; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \right) \\ &\leq C\|\phi; \dot{H}_2^{n/2-2}\| + C\|v; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\|, \\ \|\Psi(u, v); \mathcal{X}_0\| &\leq C\|\psi; L^2\| + C\|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \|u; L^2(\mathbb{R}; L^{2^*})\|, \\ \|\Psi(u, v); \dot{\mathcal{X}}\| &\leq C\|\psi; \dot{H}_2^{n/2-2}\| + C\|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \|u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\|. \end{aligned}$$

Similarly, we estimate $P_{\delta, m}\Phi(u, v)$ and $P_{2\delta, 2m}\Psi(u, v)$ in $\dot{\mathcal{X}}$

$$\begin{aligned} \|P_{\delta, m}\Phi(u, v); \dot{\mathcal{X}}\| &\leq C\|e^{\delta \cdot x} \phi; \dot{H}_2^{n/2-2}\| + C\|P_{2\delta, 2m} v; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\| \|P_{-\delta, m} u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\|, \\ \|P_{2\delta, 2m}\Psi(u, v); \dot{\mathcal{X}}\| &\leq C\|e^{2\delta \cdot x} \psi; \dot{H}_2^{n/2-2}\| + C\|P_{\delta, m} u; L^2(\mathbb{R}; \dot{B}_{2^*}^{n/2-2})\|^2, \end{aligned}$$

where C is independent of δ . We take supremum over $\delta \in D$ on both sides of the above inequalities to obtain

$$\begin{aligned} \sup_{\delta \in D} \|P_{\delta,m}\Phi(u,v);\dot{\mathcal{X}}\| &\leq C \sup_{\delta \in D} \|e^{\delta \cdot x} \phi; \dot{H}_2^{n/2-2}\| + C \sup_{\delta \in D} \|P_{2\delta,2m}v;\dot{\mathcal{X}}\| \sup_{\delta \in D} \|P_{-\delta,m}u;\dot{\mathcal{X}}\|, \\ \sup_{\delta \in D} \|P_{2\delta,2m}\Psi(u,v);\dot{\mathcal{X}}\| &\leq C \sup_{\delta \in D} \|e^{2\delta \cdot x} \psi; \dot{H}_2^{n/2-2}\| + C \left(\sup_{\delta \in D} \|P_{\delta,m}u;\dot{\mathcal{X}}\| \right)^2. \end{aligned}$$

For $(u,v), (u',v') \in G_1^D(J_m) \times G_2^D(J_{2m})$, similarly we estimate

$$\begin{aligned} \|\Phi(u,v) - \Phi(u',v'); \mathcal{X}_0\| &\leq C(\|u;\dot{\mathcal{X}}\| + \|v';\dot{\mathcal{X}}\|) \left(\|u - u'; \mathcal{X}_0\| + \|v - v'; \mathcal{X}_0\| \right), \\ \|\Psi(u,v) - \Psi(u',v'); \mathcal{X}_0\| &\leq C(\|u;\dot{\mathcal{X}}\| + \|u';\dot{\mathcal{X}}\|) \left(\|u - u'; \mathcal{X}_0\| + \|v - v'; \mathcal{X}_0\| \right), \\ \|\Phi(u,v) - \Phi(u',v'); \dot{G}_1^D(J_m)\| \\ &\leq C(\|u;\dot{G}_1^D(J_m)\| + \|v';\dot{G}_2^D(J_{2m})\|) \left(\|u - u'; \dot{G}_1^D(J_m)\| + \|v - v'; \dot{G}_2^D(J_{2m})\| \right), \\ \|\Psi(u,v) - \Psi(u',v'); \dot{G}_2^D(J_{2m})\| \\ &\leq C(\|u;\dot{G}_1^D(J_m)\| + \|u';\dot{G}_1^D(J_m)\|) \left(\|u - u'; \dot{G}_1^D(J_m)\| + \|v - v'; \dot{G}_2^D(J_{2m})\| \right). \end{aligned}$$

For $n = 3$, by Lemmas 2.2 and 2.4 and the Hölder inequality on Lorentz spaces with two exponents $3/4 = 1/2 + 1/\infty + 1/4$, $1/2 = 1/\infty + 1/\infty + 1/2$, we estimate

$$\begin{aligned} \|\Phi(u,v);\mathcal{Y}_0\| &\leq C\|\phi;L^2\| + C\|v\bar{u};L^{4/3,2}(\mathbb{R};L^{3/2})\| \\ &\leq C\|\phi;L^2\| + C\left\| |t|^{-1/2} \left\| |J_{2m}|^{1/2}(t)v(t);L^2 \right\| \left\| u(t);L^3 \right\|; L_t^{4/3,2} \right\| \\ &\leq C\|\phi;L^2\| + C\left\| |t|^{-1/2};L_t^{2,\infty} \right\| \left\| |J_{2m}|^{1/2}v;L^\infty(\mathbb{R};L^2) \right\| \left\| u;L^{4,2}(\mathbb{R};L^3) \right\| \\ &\leq C\|\phi;L^2\| + C\left\| |J_{2m}|^{1/2}v;L^\infty(\mathbb{R};L^2) \right\| \left\| u;L^{4,2}(\mathbb{R};L^3) \right\| \\ &\leq C\|\phi;L^2\| + C\|v;\dot{\mathcal{Y}}_{2m}\| \|u;\mathcal{Y}_0\|, \end{aligned}$$

$$\begin{aligned} \|\Phi(u,v);\dot{\mathcal{Y}}_m\| &\leq C\|\phi;\dot{L}_{1/2}^2\| + C\left\| |J_m|^{1/2}v\bar{u};L^{4/3,2}(\mathbb{R};L^{3/2}) \right\| \\ &\leq C\|\phi;\dot{L}_{1/2}^{1/2}\| + C\left\| |t|^{-1/2} \left\| |J_m|^{1/2}(t)u(t);L^2 \right\| \left\| |J_{2m}|^{1/2}(t)v(t);L^3 \right\| \right. \\ &\quad \left. + |t|^{-1/2} \left\| |J_m|^{1/2}(t)u(t);L^3 \right\| \left\| |J_{2m}|^{1/2}(t)v(t);L^2 \right\|; L_t^{4/3,2} \right\| \\ &\leq C\|\phi;\dot{L}_{1/2}^2\| + C\left(\left\| |J_m|^{1/2}u;L^\infty(\mathbb{R};L^2) \right\| \left\| |J_{2m}|^{1/2}v;L^{4,2}(\mathbb{R};L^3) \right\| \right. \\ &\quad \left. + \left\| |J_m|^{1/2}u;L^{4,2}(\mathbb{R};L^3) \right\| \left\| |J_{2m}|^{1/2}v;L^\infty(\mathbb{R};L^2) \right\| \right) \\ &\leq C\|\phi;\dot{L}_{1/2}^2\| + C\|v;\dot{\mathcal{Y}}_{2m}\| \|u;\dot{\mathcal{Y}}_m\|, \end{aligned}$$

and we also have

$$\begin{aligned} \|\Psi(u,v);\mathcal{Y}_0\| &\leq C\|\psi;L^2\| + C\|u;\dot{\mathcal{Y}}_m\| \|u;\mathcal{Y}_0\|, \\ \|\Psi(u,v);\dot{\mathcal{Y}}_{2m}\| &\leq C\|\psi;\dot{L}_{1/2}^2\| + C\|u;\dot{\mathcal{Y}}_m\|^2. \end{aligned}$$

Similarly, we estimate $P_{\delta,m}\Phi(u, v)$ and $P_{2\delta,2m}\Psi(u, v)$ in $\mathcal{Y}_0, \dot{\mathcal{Y}}_m$ and $\dot{\mathcal{Y}}_{2m}$ as follows

$$\begin{aligned} \|P_{\delta,m}\Phi(u, v); \mathcal{Y}_0\| &\leq C\|e^{\delta \cdot x}\phi; L^2\| + C\|P_{2\delta,2m}v; \dot{\mathcal{Y}}_{2m}\|\|P_{-\delta,m}u; \mathcal{Y}_0\|, \\ \|P_{2\delta,2m}\Psi(u, v); \mathcal{Y}_0\| &\leq C\|e^{2\delta \cdot x}\psi; L^2\| + C\|P_{\delta,m}u; \dot{\mathcal{Y}}_m\|\|P_{\delta,m}u; \mathcal{Y}_0\|, \end{aligned}$$

and

$$\begin{aligned} \|P_{\delta,m}\Phi(u, v); \dot{\mathcal{Y}}_m\| &\leq C\|e^{\delta \cdot x}\phi; \dot{L}_{1/2}^2\| + C\|P_{2\delta,2m}v; \dot{\mathcal{Y}}_{2m}\|\|P_{-\delta,m}u; \dot{\mathcal{Y}}_m\|, \\ \|P_{2\delta,2m}\Psi(u, v); \dot{\mathcal{Y}}_{2m}\| &\leq C\|e^{2\delta \cdot x}\psi; \dot{L}_{1/2}^2\| + C\|P_{\delta,m}u; \dot{\mathcal{Y}}_m\|^2, \end{aligned}$$

where C is independent of δ . We take supremum over $\delta \in D$ on both sides of the above inequalities to obtain

$$\begin{aligned} \sup_{\delta \in D} \|P_{\delta,m}\Phi(u, v); \mathcal{Y}_0\| &\leq C \sup_{\delta \in D} \|e^{\delta \cdot x}\phi; L^2\| + C \sup_{\delta \in D} \|P_{2\delta,2m}v; \dot{\mathcal{Y}}_{2m}\| \sup_{\delta \in D} \|P_{-\delta,m}u; \mathcal{Y}_0\|, \\ \sup_{\delta \in D} \|P_{2\delta,2m}\Psi(u, v); \mathcal{Y}_0\| &\leq C \sup_{\delta \in D} \|e^{2\delta \cdot x}\psi; L^2\| + C \sup_{\delta \in D} \|P_{\delta,m}u; \dot{\mathcal{Y}}_m\| \sup_{\delta \in D} \|P_{\delta,m}u; \mathcal{Y}_0\|, \\ \sup_{\delta \in D} \|P_{\delta,m}\Phi(u, v); \dot{\mathcal{Y}}_m\| &\leq C \sup_{\delta \in D} \|e^{\delta \cdot x}\phi; \dot{L}_{1/2}^2\| + C \sup_{\delta \in D} \|P_{2\delta,2m}v; \dot{\mathcal{Y}}_{2m}\| \sup_{\delta \in D} \|P_{-\delta,m}u; \dot{\mathcal{Y}}_m\|, \\ \sup_{\delta \in D} \|P_{2\delta,2m}\Psi(u, v); \dot{\mathcal{Y}}_{2m}\| &\leq C \sup_{\delta \in D} \|e^{2\delta \cdot x}\psi; \dot{L}_{1/2}^2\| + C \left(\sup_{\delta \in D} \|P_{\delta,m}u; \dot{\mathcal{Y}}_m\| \right)^2. \end{aligned}$$

For $(u, v), (u', v') \in G_1^D(J_m) \times G_2^D(J_{2m})$, similarly we estimate

$$\begin{aligned} &\|\Phi(u, v) - \Phi(u', v'); G_{1,0}^D(J_m)\| \\ &\leq C(\|u; \dot{G}_m^D(J_m)\| + \|v'; \dot{G}_{2m}^D(J_{2m})\|) \left(\|u - u'; G_{1,0}^D(J_m)\| + \|v - v'; G_{2,0}^D(J_{2m})\| \right), \\ &\|\Psi(u, v) - \Psi(u', v'); G_{2,0}^D(J_{2m})\| \\ &\leq C(\|u; \dot{G}_m^D(J_m)\| + \|u'; \dot{G}_m^D(J_m)\|) \left(\|u - u'; G_{1,0}^D(J_m)\| + \|v - v'; G_{2,0}^D(J_{2m})\| \right), \\ &\|\Phi(u, v) - \Phi(u', v'); \dot{G}_1^D(J_m)\| \\ &\leq C(\|u; \dot{G}_1^D(J_m)\| + \|v'; \dot{G}_2^D(J_{2m})\|) \left(\|u - u'; \dot{G}_1^D(J_m)\| + \|v - v'; \dot{G}_2^D(J_{2m})\| \right), \\ &\|\Psi(u, v) - \Psi(u', v'); \dot{G}_2^D(J_{2m})\| \\ &\leq C(\|u; \dot{G}_1^D(J_m)\| + \|u'; \dot{G}_1^D(J_m)\|) \left(\|u - u'; \dot{G}_1^D(J_m)\| + \|v - v'; \dot{G}_2^D(J_{2m})\| \right). \end{aligned}$$

Therefore for any $(u, v), (u', v') \in X(R, \varepsilon)$ with $n \geq 4$, we have

$$\begin{aligned} &\left\| \left(\Phi(u, v), \Psi(u, v) \right); \mathcal{X}_0 \times \mathcal{X}_0 \right\| \leq C\|(\psi, \phi); L^2 \times L^2\| + CR\varepsilon, \\ &\left\| \left(\Phi(u, v), \Psi(u, v) \right); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m}) \right\| \leq C\rho + CR\varepsilon, \\ &\left\| \left(\Phi(u, v) - \Phi(u', v'), \Psi(u, v) - \Psi(u', v') \right); \mathcal{X}_0 \times \mathcal{X}_0 \right\| \leq C\varepsilon\|(u - u', v - v'); \mathcal{X}_0 \times \mathcal{X}_0\|, \\ &\left\| \left(\Phi(u, v) - \Phi(u', v'), \Psi(u, v) - \Psi(u', v') \right); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m}) \right\| \\ &\leq C\varepsilon\|(u - u', v - v'); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m})\|, \end{aligned}$$

and with $n = 3$, we have

$$\begin{aligned}
& \left\| \left(\Phi(u, v), \Psi(u, v) \right); G_{1,0}^D(J_m) \times G_{2,0}^D(J_{2m}) \right\| \leq C \|(\phi, \psi); G_1^D(x; L^2) \times G_2^D(x; L^2)\| + CR\varepsilon, \\
& \left\| \left(\Phi(u, v), \Psi(u, v) \right); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m}) \right\| \leq C\rho + CR\varepsilon, \\
& \left\| \left(\Phi(u, v) - \Phi(u', v'), \Psi(u, v) - \Psi(u', v') \right); G_{1,0}^D(J_m) \times G_{2,0}^D(J_{2m}) \right\| \\
& \leq C\varepsilon \| (u - u', v - v'); G_{1,0}^D(J_m) \times G_{2,0}^D(J_{2m}) \|, \\
& \left\| \left(\Phi(u, v) - \Phi(u', v'), \Psi(u, v) - \Psi(u', v') \right); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m}) \right\| \\
& \leq C\varepsilon \| (u - u', v - v'); \dot{G}_1^D(J_m) \times \dot{G}_2^D(J_{2m}) \|.
\end{aligned}$$

Therefore for any $(\phi, \psi) \in B_1^D(\rho) \times B_2^D(\rho)$ with R, ε and ρ satisfying

$$\begin{cases} C \|(\phi, \psi); L^2 \times L^2\| + CR\varepsilon \leq R, \\ C\rho + CR\varepsilon \leq \varepsilon, \\ C\varepsilon < 1, \end{cases}$$

for $n \geq 4$

$$\begin{cases} C \|(\phi, \psi); G_1^D(x; L^2) \times G_2^D(x; L^2)\| + CR\varepsilon \leq R, \\ C\rho + CR\varepsilon \leq \varepsilon, \\ C\varepsilon < 1, \end{cases}$$

for $n = 3$, the mapping $(\Phi, \Psi) : (u, v) \mapsto (\Phi(u, v), \Psi(u, v))$ is a contraction in $G_1^D(J_m) \times G_2^D(J_{2m})$ and has a unique fixed point in $G_1^D(J_m) \times G_2^D(J_{2m})$. This completes the proof of Theorem 1.1.

References

- [1] J.-L. Bona, Z. Grujić and H. Kalisch, Global solutions of the derivative Schrödinger equation in a class of functions analytic in a strip, *J. Differential Equations.*, **229**,(2006), 186-203.
- [2] J.-M. Bony, Calcul symbolique et propagation des singularité pour les équations aux dérivées partielles non linéaires, *Ann. Sci. École Norm. Sup.*, (4) **14**, (1981), 209-246.
- [3] T. Cazenave, Equation de Schrödinger non linéaires en dimension deux, *Proc. Roy. Soc. Edinburgh Sect. A.*, **84**, (1979), 327-346.
- [4] T. Cazenave, Semilinear Schrödinger Equations, *Courant Lecture Notes in Math.*, 10, Amer. Math. Soc., 2003.
- [5] T. Cazenave, and F. B. Weissler, The Cauchy problem for the critical nonlinear Schrödinger equation in H^s , *Nonlinear Analysis, TMA.*, **14**, (1990), 807-836.
- [6] A. DeBouard, Analytic solution to non elliptic non linear Schrödinger equations, *J. Differential Equations.*, **104**, (1993), 196-213.
- [7] J. Ginibre, Introduction aux équations de Schrödinger non linéaires, *Paris Onze Edition, L161. Université Paris-Sud*, 1998.
- [8] N. Hayashi, Global existence of small analytic solutions to nonlinear Schrödinger equations, *Duke Math. J.*, **60**, (1990), 717-727.
- [9] N. Hayashi, Smoothing effect of small analytic solutions to nonlinear Schrödinger equations, *Ann. Inst. H. Poincaré, Phys. Théor.*, **57**, (1992), 717-727.
- [10] N. Hayashi and K. Kato, Analyticity in time and smoothing effect of solutions to nonlinear Schrödinger equations, *Commun. Math. Phys.*, **184**, (1997), 273-300.
- [11] N. Hayashi, C. Li and P. I. Naumkin, On a system of nonlinear Schrödinger equations in 2d, *Diff. Int. Eqs.*, **24**, (2011), 417-434.
- [12] N. Hayashi, C. Li and P. I. Naumkin, Modified wave operator for a system of nonlinear Schrödinger equations in 2d, *Commun. Partial Differential Equations.*, **37**, (2012), 947-968.
- [13] N. Hayashi, C. Li and T. Ozawa, Small data scattering for a system of nonlinear Schrödinger equations, *Differ. Equ. Appl.*, **3**, (2011), 415-426.
- [14] N. Hayashi and T. Ozawa, Scattering theory in the weighted $L^2(\mathbb{R}^n)$ spaces for some nonlinear Schrödinger equations, *Ann. Inst. H. Poincaré Phys. Théor.*, **48**, (1988), 17-37.
- [15] N. Hayashi and T. Ozawa, On the derivative nonlinear Schrödinger equation, *Physica D*, **55**, (1992), 14-36.
- [16] N. Hayashi T. Ozawa and K. Tanaka, On a system of nonlinear Schrödinger equations with quadratic interaction, *Ann. Inst. Henri Poincaré, Analyse non linéaire*, **30**, (2013), 661-690.
- [17] N. Hayashi and S. Saitoh, Analyticity and smoothing effect for the Schrödinger equation, *Ann. Inst. Henri Poincaré, Phys. Théor.*, **52**, (1990), 163-173.
- [18] N. Hayashi and S. Saitoh, Analyticity and global existence of small solutions to some nonlinear Schrödinger equations, *Commun. Math. Phys.*, **129**, (1990), 27-41.
- [19] G. Hoshino and T. Ozawa, Analytic smoothing effect for nonlinear Schrödinger equation in two space dimensions, *Osaka J. Math.*, (in press).
- [20] G. Hoshino and T. Ozawa, Analytic smoothing effect for a system of nonlinear Schrödinger equations, *Differ. Equ. Appl.*, **5**, (2013), 395-408.
- [21] G. Hoshino and T. Ozawa, Analytic smoothing effect for nonlinear Schrödinger equation with quintic nonlinearity, (submitted).

- [22] M. Keel and T. Tao, Endpoint Strichartz inequalities, *Amer. J. Math.*, **120**, (1998), 955-980.
- [23] K. Kato and K. Taniguchi, Gevrey regularizing effect for nonlinear Schrödinger equations, *Osaka. J. Math.*, **33**, (1996), 863-880.
- [24] T. Kato, On nonlinear Schrödinger equations. II H^s -solutions and unconditional well-posedness, *J. d'Anal. Math.*, **67**, (1995), 281-306.
- [25] T. Kato and K. Masuda, Nonlinear evolution equations and analyticity, *Ann. Inst. Henri Poincaré, Analyse non linéaire.*, **3**, (1986), 455-467.
- [26] H. Kozono, T. Ogawa and Y. Taniuchi, Navier-Stokes equations in the Besov space near L^∞ and BMO , *Kyushu. J. Math.*, **57**, (2003), 303-324.
- [27] K. Nakamitsu, Analytic finite energy solutions of the nonlinear Schrödinger equation, *Commun. Math. Phys.*, **260**, (2005), 117-130.
- [28] M. Nakamura and T. Ozawa, Low energy scattering for nonlinear Schrödinger equations in fractional order Sobolev spaces, *Rev. Math. Phys.*, **9**, (1997), 397-410.
- [29] M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, *J. Funct. Anal.*, **155**, (1998), 364-380.
- [30] K. Nakanishi, Asymptotically-free solutions for the short range nonlinear Schrödinger equation, *SIAM J. Math. Anal.*, **32**, (2001), 1265-1271.
- [31] K. Nakanishi and T. Ozawa, Remarks on scattering for nonlinear Schrödinger equations, *NODEA Nonlinear Differ. Equ. Appl.*, **9**, (2002), 45-68.
- [32] T. Ozawa and K. Yamauchi, Remarks on analytic smoothing effect for the Schrödinger equation, *Math. Z.*, **261**, (2009), 511-524.
- [33] T. Ozawa and K. Yamauchi, Analytic smoothing effect for global solutions to nonlinear Schrödinger equation, *J. Math. Anal Appl.*, **364**, (2010), 492-497.
- [34] T. Ozawa, K. Yamauchi and Y. Yamazaki, Analytic smoothing effect for solutions to Schrödinger equations with nonlinearity of integral type, *Osaka J. Math.*, **42**, (2005), 737-750.
- [35] M. Reed and B. Simon, Method of Mathematical Physics II, *Academic Press*, 1975.
- [36] L. Robbiano and C. Zuily, Microlocal analytic smoothing effect for the Schrödinger equation, *Duke Math. J.*, **100**, (1999), 93-129.
- [37] L. Robbiano and C. Zuily, Effect régularisant microlocal analytique pour l'équation de Schrödinger : le cas données oscillantes, *Comm. Partial Differential Equations*, **25**, (2000), 1891-1906.
- [38] R. S. Strichartz, Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations, *Duke. Math. J.*, **44**, (1977), 705-714.
- [39] J. C. H Simon and E. Taflin, Wave operators and analytic solutions of nonlinear Klein-Gordon equations and of nonlinear Schrödinger equations, *Commun. Math. Phys.*, **99**, (1985), 541-562.
- [40] E. M. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, *Princeton Univ. Press*, 1971.
- [41] E. M. Stein, G. Weiss and M. Weiss, H^p classes of holomorphic functions in tube domains, *Proc. N. A. S.*, **52** (1964), 1035-1039.
- [42] C. Sulem and P.-L. Sulem, The Nonlinear Schrödinger Equation. Self-focusing and Wave Collapse, *Appl. Math. Sci.*, **139**, Springer 1999.
- [43] H. Takuwa, Analytic smoothing effects for a class of dispersive equations, *Tsukuba J. Math.*, **28**, (2004), 1-34.
- [44] K. Yajima, Existence of solutions for Schrödinger evolution equations, *Commun. Math.*

Phys., **110**, (1987), 415-426.